

Geometric creation of quantum vorticity

Michael R.R. Good^{a,b} Chi Xiong^b Alvin J.K. Chua^{b,c} Kerson Huang^{b,d}

^a*Department of Physics, Nazarbayev University, Astana, Kazakhstan*

^b*Institute of Advanced Studies, Nanyang Technological University, Singapore*

^c*Institute of Astronomy, University of Cambridge, Cambridge, UK*

^d*Department of Physics, Massachusetts Institute of Technology, Cambridge, MA, USA*

E-mail: michael.good@nu.edu.kz, xiongchi@ntu.edu.sg,
ajkc3@ast.cam.ac.uk, kerson@mit.edu

ABSTRACT: We consider superfluidity and quantum vorticity in rotating spacetimes. The system is described by a complex scalar satisfying a nonlinear Klein-Gordon equation. Rotation terms are identified and found to lead to the transfer of angular momentum of the spacetime to the scalar field. The scalar field responds by rotating, physically behaving as a superfluid, through the creation of quantized vortices. We demonstrate the vortex nucleation through numerical simulation.

Keywords: quantum vorticity, Kerr metric, BTZ metric

Contents

1	Introduction	1
2	NLKG in an arbitrary metric	2
3	NLKG in the BTZ metric	3
4	NLKG in the Kerr metric	6
5	Conclusions and Outlooks	8

1 Introduction

It is known that the vacuum is not empty but filled with at least one complex scalar field, namely the Higgs field postulated to generate mass in the Standard Model of particle physics. Its existence has found experimental support in the discovery of the associated field quanta, the Higgs boson [1, 2]. The nonlinear quantum phase dynamics of such a field, on a macroscopic scale, can lead to superfluidity. There are proposals that associate many cosmological phenomena with this superfluidity, chief among which are dark energy, dark matter, and inflation [3]-[14].

In this paper, we study the superfluidity of the Higgs vacuum in rotating spacetimes. Quantized vorticity is the only means through which a superfluid can rotate, and as such, stands as a prominent signature of superfluidity. Steady regularly-distributed vortex lattices have been experimentally observed in rotating superfluid helium [15] and the rotating Bose-Einstein condensates (BEC) of cold trapped atoms [16]. This is a very interesting and distinctive macroscopic quantum phenomenon, and one may wonder whether it can happen on astrophysical or cosmological scales. Our observation is that rotating black holes, such as the Kerr black hole in (3+1)-dimensions [17] and the BTZ black hole in (2+1)-dimensions [18], can provide a similar rotating environment as in condensed physics experiments and hence may induce quantum vortices in the cosmic superfluid or the Higgs vacuum. However, the rotating “bucket” here, significantly different from the container for liquid helium and the laser trap for BECs, is the spacetime itself. It has never been demonstrated in principle that curved spacetime itself can nucleate a quantum vortex lattice.

In general relativity, frame-dragging (or the Lense-Thirring effect [19]) refers to a special distortion of spacetime geometry caused by a rotating mass. It has been detected [20], and occurs because the spacetime has angular momentum. We explore the possibility that the frame-dragging effect of a spacetime with angular momentum can create quantum vortices in the cosmic superfluid. This is quantum vortex formation induced by angular momentum transfer from the spacetime to the field. We refer to it as *geometrical creation of quantized vorticity*.

In both the superfluid helium and the BEC cases, the system is described by a macroscopic wave function $\Psi(\vec{x}, t) = |\Psi(\vec{x}, t)|e^{i\alpha(\vec{x}, t)}$ as an order parameter. $\Psi(\vec{x}, t)$ satisfies the nonlinear Schrödinger equation (NLSE) or the Gross-Pitaevskii equation (GPE), which are not suitable for relativistic systems. Therefore we use the nonlinear Klein-Gordon equation (NLKG), promoting the order parameter $\Psi(\vec{x}, t)$ to a complex scalar field $\Phi(\vec{x}, t)$. We include a usual nonlinear potential (for simplicity we use a Higgs potential in the present paper), but the main difference here is the use of the non-trivial topology of the curved space NLKG equation, which will be subject to the effect of black hole spacetimes.

In general, this Klein-Gordon equation should be solved in combination with the Einstein equation; however, we neglect the backreaction of the complex scalar field $\Phi(\vec{x}, t)$ on the stress-energy tensor, and treat the rotating black hole as the spacetime background in the canonical semi-classical approach. Despite neglecting backreaction, it is difficult to find numerical solutions for the NLKG in such backgrounds. We are therefore compelled to make a slow-rotation approximation under which two rotation terms, a Coriolis term and a centrifugal term, emerge at first and second order in the rotational angular velocity Ω , respectively. Based on numerical computations, we find that the first order Coriolis term is the most important. We show that it leads to vortex formation, and therefore that quantum vortices can be generated by the frame-dragging effect of black holes. The fast-rotating case is left for future investigation.

An interesting finding is that the emergence of the rotation terms occurs for both the BTZ case and the Kerr case. We show this explicitly by expanding their equations of motion in terms of angular velocity. We also find a direct link between the BTZ background and a simple rotation metric. We numerically evolve the field in the (2+1)-dimensional case with appropriate approximations, looking for quantum vorticity. Vortex nucleation in this case emerges in a similar way to superfluid helium and BEC vortex lattices. We have not solved the more formidable Kerr case. Even though we have identified the same indispensable rotation terms in the Kerr equations of motion, the Kerr background geometry presents a significantly more challenging numerical endeavor. Nevertheless, in the Kerr case (like the BTZ case), it is still certain that the Coriolis rotation term will be responsible for the transfer of angular momentum to the field, catalyzing quantum vortex formation.

2 NLKG in an arbitrary metric

Setting $\hbar = 1$, we consider a generic complex scalar field Φ , not particularly tied to any vacuum field in particle theory, described phenomenologically by a nonlinear Klein-Gordon equation (NLKG)

$$\square\Phi + \lambda(|\Phi|^2 - F_0^2)\Phi = 0, \quad (2.1)$$

where

$$\square\Phi \equiv (-g)^{-1/2} \partial^\mu \left[(-g)^{1/2} g_{\mu\nu} \partial^\nu \Phi \right]. \quad (2.2)$$

The metric $g_{\mu\nu}$ is the means through which frame-dragging effects will enter, while λ and F_0 are the self-interaction constant of the Φ field and the vacuum expectation value of its modulus, respectively. The nonlinear self-interaction spontaneously breaks the global gauge

invariance, and maintains a nonzero vacuum field F_0 . As shown in [13], this vacuum field is stable against collapse due to self-gravitation, i.e., the equivalent Jeans length is greater than the radius of the universe. The NLKG equation is used (as opposed to the non-linear Schrödinger equation), because relativistic effects are essential in this context. In addition, the NLKG equation is more generalizable to quantum field theory in curved spacetime, and the relationship between angular velocity and the number of vortices (Feynman's relation) becomes modified in the relativistic regime [21].

In a phase representation $\Phi = Fe^{i\sigma}$, the superfluid velocity is given by $\mathbf{v}_s = \xi_s \nabla \sigma$, where $\xi_s = c^2 (-\partial\sigma/\partial t)^{-1}$ is a spacetime dependent factor that makes $|\mathbf{v}_s| < c$. In the non-relativistic limit we have $\xi_s \rightarrow \hbar/m$, where m is a mass scale [13, 21]. In practice, it is simpler to work with $\nabla\sigma$. A quantized vortex is a solution to the NLKG with

$$\oint_C \nabla\sigma \cdot d\mathbf{s} = 2\pi n \quad (n = 0, \pm 1, \pm 2, \dots), \quad (2.3)$$

over some spatial closed loop C . Such quantized vortices have been extensively studied in liquid helium and Bose-Einstein condensates of cold trapped atoms, both experimentally and theoretically [13, 15, 16, 21]. We will search for these quantized vortex solutions in the following section.

3 NLKG in the BTZ metric

We want to extract specific terms in the NLKG, Eq. (2.1), that arise from the metric and induce vortex creation. For orientation, we consider first the BTZ metric [18] in (2+1)-dimensional spacetime, which is a vacuum solution to Einstein's equation in the region outside of a rotating "star" (a central distribution that has both mass and angular momentum). The line element in spatial polar coordinates is given by

$$ds_B^2 = g_{tt}dt^2 + g_{rr}dr^2 + g_{\phi\phi}d\phi^2 + 2g_{t\phi}dtd\phi, \quad (3.1)$$

or

$$ds_B^2 = \left(g_{tt} - \frac{g_{t\phi}^2}{g_{\phi\phi}} \right) dt^2 + g_{rr}dr^2 + g_{\phi\phi} \left(d\phi + \frac{g_{t\phi}}{g_{\phi\phi}} dt \right)^2, \quad (3.2)$$

with components in matrix form using (t, r, ϕ) coordinates,

$$(g_{\mu\nu})^{\text{BTZ}} = \begin{pmatrix} 8GM - \frac{c^2 r^2}{a^2} & 0 & -\frac{4GJ}{c^2} \\ 0 & \frac{r^2}{a^2} + \frac{16G^2 J^2}{c^6 r^2} - \frac{8GM}{c^2} & 0 \\ -\frac{4GJ}{c^2} & 0 & r^2 \end{pmatrix}, \quad (3.3)$$

where M is the black hole mass, J is the angular momentum of the spacetime and a is the AdS radius. The BTZ black hole is a solution for (2+1)-dimensional gravity with a negative cosmological constant, i.e. asymptotically AdS. It is used as a toy model to illustrate relevant physical content.

The frame-dragging component $g_{t\phi}$ implies a local rotational angular velocity

$$\Omega_B = -\frac{g_{t\phi}}{g_{\phi\phi}} = \frac{4GJ}{c^2 r^2}, \quad (3.4)$$

which is independent of the mass M . The NLKG equation in the BTZ metric is Eq. (2.1) with $g_{\mu\nu}$ given by Eq. (3.3),¹ i.e.,

$$\begin{aligned} \square\Phi &= -g_{rr}\left(\frac{\partial^2\Phi}{\partial t^2} + \frac{J}{r^2}\frac{\partial^2\Phi}{\partial t\partial\phi} + \frac{J^2}{4r^4}\frac{\partial^2\Phi}{\partial\phi^2}\right) + \frac{1}{r^2}\frac{\partial^2\Phi}{\partial\phi^2} + \frac{1}{r}\frac{\partial}{\partial r}\left(\frac{r}{g_{rr}}\frac{\partial\Phi}{\partial r}\right) \\ &= -\lambda(|\Phi|^2 - F_0^2)\Phi, \end{aligned} \quad (3.5)$$

where g_{rr} is given by

$$g_{rr} = \frac{1}{\frac{J^2}{4r^2} + \frac{r^2}{a^2} - M} = \frac{4a^2 r^2}{4r^4 + a^2(J^2 - 4Mr^2)}. \quad (3.6)$$

To illustrate the structure of Eq. (3.5), we apply the usual method of separation of variables:

$$\Phi(t, r, \phi) = e^{-i\omega t} e^{im\phi} R(r) \quad (3.7)$$

to obtain

$$\left[g_{rr}\left(\omega - m\Omega_B\right)^2 - \frac{m^2}{r^2} + \frac{1}{r}\frac{d}{dr}\left(\frac{r}{g_{rr}}\frac{d}{dr}\right) \right] R(r) = \lambda(R^2 - F_0^2)R(r). \quad (3.8)$$

Interestingly, the corresponding homogeneous differential equation of Eq. (3.8) (i.e. removing the nonlinear potential term on the right-hand side of Eq. (3.8)) is actually a hypergeometric equation [22]. However, the ansatz Eq. (3.7) cannot lead to a vortex lattice solution and we have to solve the equation Eq. (3.5) without it.

Reinstating G and c , we now show that under an appropriate approximation, solving Eq. (3.5) is equivalent to solving the NLKG in the following comparison metric that describes rotation in flat (2+1)-dimensional spacetime, with angular velocity Ω_0 :

$$ds_{\text{comparison}}^2 = -c^2 dt^2 + dr^2 + r^2(d\phi - \Omega_0 dt)^2. \quad (3.9)$$

The effect of rotation enters the NLKG only through $\square\Phi$, which separates into a non-rotational term plus terms that can be identified with the Coriolis force and centrifugal force:

$$\square\Phi = \square^{(0)}\Phi + R_{\text{Coriolis}} + R_{\text{centrifugal}}, \quad (3.10)$$

where $\square^{(0)}\Phi$ is that for the (2+1)-dimensional Minkowski metric,

$$\square^{(0)}\Phi(t, r, \phi) = -\frac{1}{c^2}\frac{\partial^2\Phi(t, r, \phi)}{\partial t^2} + \frac{\partial^2\Phi(t, r, \phi)}{\partial r^2} + \frac{1}{r}\frac{\partial\Phi(t, r, \phi)}{\partial r} + \frac{1}{r^2}\frac{\partial^2\Phi(t, r, \phi)}{\partial\phi^2}, \quad (3.11)$$

¹Here we temporarily take the units that are usual for the BTZ metric, $c = 1$ and $G = 1/8$, to illustrate the simplicity of the hypergeometric result.

and the rotational terms are

$$R_{\text{Coriolis}} = -\frac{2\Omega_0}{c^2} \frac{\partial^2 \Phi}{\partial t \partial \phi}, \quad R_{\text{centrifugal}} = -\frac{\Omega_0^2}{c^2} \frac{\partial^2 \Phi}{\partial \phi^2}. \quad (3.12)$$

The equivalence between the two approaches comes from the fact that the BTZ metric is formally connected to the comparison metric for the special values

$$M = -\frac{c^2}{8G}, \quad a^2 = -\frac{c^2}{\Omega_0^2}, \quad (3.13)$$

which leads to

$$g_{rr} = \frac{1}{1 + \frac{r^2}{c^2}(\Omega_B^2 - \Omega_0^2)} = 1 - \frac{r^2}{c^2}(\Omega_B^2 - \Omega_0^2) + \mathcal{O}((\Omega_B^2 - \Omega_0^2)^2), \quad (3.14)$$

and hence

$$(g_{\mu\nu})^{\text{BTZ}} - (g_{\mu\nu})^{\text{Comparison}} = \begin{pmatrix} 0 & 0 & (\Omega_0 - \Omega_B)r^2 \\ 0 & -\frac{r^2}{c^2}(\Omega_B^2 - \Omega_0^2) & 0 \\ (\Omega_0 - \Omega_B)r^2 & 0 & 0 \end{pmatrix} + \mathcal{O}((\Omega_B^2 - \Omega_0^2)^2). \quad (3.15)$$

Therefore the difference between these two metrics vanishes if we assume that Ω_B is approximately constant: $\Omega_B \approx \Omega_0$. Under this approximation, it is easy to read Eq. (3.10), Eq. (3.11) and Eq. (3.12) from Eq. (3.5). (Alternatively, one can simply write the NLKG in the comparison metric to identify the terms R_{Coriolis} and $R_{\text{centrifugal}}$ in Eq. (3.12) [21].)

In general, we can identify the effective Coriolis force and centrifugal force arising from the BTZ metric by expanding Eq. (3.5) in powers of J (or equivalently Ω_B) to second order, which gives:

$$R_{\text{Coriolis}}^B = \frac{2\Omega_B}{g_{tt}} \frac{\partial^2 \Phi}{\partial t \partial \phi}, \quad R_{\text{centrifugal}}^B = \frac{\Omega_B^2}{g_{tt}} \frac{\partial^2 \Phi}{\partial \phi^2}. \quad (3.16)$$

There are of course higher order terms in the BTZ wave operator $\square\Phi$; however, the presence of the terms Eq. (3.16) indicates that the field Φ does, in fact, feel the rotation of the star through spacetime geometry. This is to be expected, since frame-dragging is spacetime dependent. The terms in Eq. (3.16) depend on r because Ω_B and g_{tt} depend on r , and they approach the form Eq. (3.12) when $g_{tt} \rightarrow -c^2$, in the limits:

$$(g_{tt})^{\text{BTZ}} \Big|_{a \rightarrow \infty}^{M = -\frac{c^2}{8G}} = -c^2, \quad (g_{tt})^{\text{BTZ}} \Big|_{r \rightarrow a}^{M \rightarrow 0} = -c^2. \quad (3.17)$$

It is noted that in the first limit of Eq. (3.17), (after M is set to its negative value), $a \rightarrow \infty$ is equivalent to the regime $r\Omega \ll c$ in the context of the formal reduction $a^2 = -c^2/\Omega^2$. In addition, when this first limit of Eq. (3.17) and $J \rightarrow 0$ is applied to the BTZ metric Eq. (3.3), the result is the flat and non-rotating (2+1)-dimensional Minkowski metric. The second limit of Eq. (3.17), where $M \rightarrow 0$, allows one to obtain the empty space vacuum state. The physical significance of the first limit lies in its consistency of approach to flat spacetime. The physical significance of the second limit lies in the fact that it furnishes

a vanishing, non-negative black hole mass, approaching the vacuum state of empty space (not AdS space). In the context of AdS space emerging as a bound state from a continuous black hole mass spectrum (after one has set $M = -c^2/8G$ and $J = 0$ but before one has taken the limit $a \rightarrow \infty$) [18], the first limit can be considered a more natural choice.

Although the rotational effects are contained in $\square\Phi$, the nonlinear self-interaction will be necessary to produce vortices. Without the nonlinear potential in Eq. (2.1), the situation is that of a free test particle in the spacetime. To see this, consider the radial field equation Eq. (3.8). The radial field describes some material in the spacetime, while the nonlinear potential brings in self-interaction. The situation here is very similar to that for the NLKG with a rotating star as the source (discussed in [13, 21]); however, the numerical computation in the BTZ case is more challenging, owing to its curved-space nature and the cross derivative in the Coriolis force.

To explicitly demonstrate geometric vortex creation, we solve the NLKG numerically in the comparison spacetime, (i.e. Eq. (2.1) with the metric Eq. (3.9)), which retains the frame-dragging rotational terms Eq. (3.12). The equivalence to the BTZ metric can be seen from Eq. (3.15). This approach is possible to handle numerically and the results can be seen in Fig. (1). For various choices of model parameters (which we specify), a vortex lattice emerges in the superfluid.

For a finite region in space whose size is very small compared with the AdS radius of the BTZ metric, the angular velocity Ω_B can be considered as approximately constant, Ω_0 . The numerical results for the NLKG in the comparison metric are equivalent to those for the NLKG in the BTZ metric under such an approximation. In Fig. (1) we used a finite-difference scheme and spectrum methods to decouple the cross derivatives ∂_t and ∂_ϕ in the Coriolis term, and a semi-implicit scheme similar to the Crank-Nicolson method for the linear terms and an explicit scheme for the nonlinear term. The boundary of the computation domain is chosen to satisfy $\Omega r \ll c$, and a second order Neumann boundary condition is imposed.

Fig. (1) contains the vortex lattice with a plot of the modulus of the field $|\Phi|$ on the left side and a plot of the phase σ on the right side. The locations of vortices are indicated by the dots (i.e. field zeros) in the modulus plot, and by the phase discontinuities around which the color changes from blue to red (see online color version), corresponding to a phase change of 2π . These results demonstrate quantum vortices. Their distribution is determined by the Coriolis term. As the computation is performed under the slow-rotation approximation $\Omega \ll 1$, the centrifugal term of order Ω^2 causes only a small non-uniformity to the vortex lattice.

4 NLKG in the Kerr metric

It is now interesting to turn to the Kerr metric² in (3+1)-dimensional space-time, which describes the region outside of a rotating star of mass M and angular momentum J [17].

²The Kerr metric may also be parametrized by the Schwarzschild surface gravity g and spring constant $k \equiv M\Omega_K^2$, in order to easily demonstrate its Hawking temperature $2\pi T = g - k$; see [23].




Figure 1. Left panel: Lattice of 100 quantized vortices. This is a contour plot of the modulus of a complex scalar field that satisfies the nonlinear Klein-Gordon equation in (2+1)-dimensional curved space-time, with frame-dragging. Right panel: Contour plot of the phase of the complex scalar field, showing the “strings” across which the phase jumps by 2π . The computation is performed in a circular domain with radius $R = 120$, and the normalized parameters are $\Omega_0 = 1/240$, $\lambda = 6.4$ and $F_0 = 1$.

The Kerr metric reduces to the Schwarzschild metric when $J = 0$. The line element is, in spatial spherical coordinates,

$$ds_K^2 = g_{tt}dt^2 + g_{rr}dr^2 + g_{\theta\theta}d\theta^2 + g_{\phi\phi}d\phi^2 + 2g_{t\phi}dtd\phi, \quad (4.1)$$

with

$$\begin{aligned} g_{tt} &= -c^2 (\Delta - \alpha^2 \sin^2 \theta) \Sigma^{-1}, & g_{rr} &= \Sigma \Delta^{-1}, \\ g_{\theta\theta} &= \Sigma, & g_{\phi\phi} &= -\sin^2 \theta \left[\alpha^2 \Delta \sin^2 \theta - (r^2 + \alpha^2)^2 \right] \Sigma^{-1}, \\ g_{t\phi} &= -c\alpha \sin^2 \theta (r^2 + \alpha^2 - \Delta) \Sigma^{-1}, \end{aligned} \quad (4.2)$$

where

$$\begin{aligned} \Delta &\equiv \alpha^2 + r^2 - rr_s, & \Sigma &\equiv \alpha^2 \cos^2 \theta + r^2, \\ r_s &\equiv 2GMc^{-2}, & \alpha &\equiv J/Mc. \end{aligned} \quad (4.3)$$

The local angular velocity is given by [24]

$$\Omega_K = -\frac{g_{t\phi}}{g_{\phi\phi}} = \frac{\alpha(r^2 + \alpha^2 - \Delta)c}{(r^2 + \alpha^2)^2 - \Delta\alpha^2 \sin^2 \theta}. \quad (4.4)$$

For a comparison metric in (3+1) dimensions, the Coriolis and centrifugal terms have the same form as in Eq. (3.12), except that ϕ is to be identified with the azimuthal angle in three spatial dimensions. We calculate $\square\Phi$ using the Kerr metric and write the NLKG Eq. (2.1) explicitly as

$$\Sigma \square\Phi = \left[-\frac{(r^2 + \alpha^2)^2}{\Delta} + \alpha^2 \sin^2 \theta \right] \frac{\partial^2 \Phi}{\partial t^2} - \frac{4\alpha Mr}{\Delta} \frac{\partial^2 \Phi}{\partial t \partial \phi} + \frac{\Sigma - 2Mr}{\Delta \sin^2 \theta} \frac{\partial^2 \Phi}{\partial \phi^2}$$

$$\begin{aligned}
& + \frac{\partial}{\partial r} \left(\Delta \frac{\partial \Phi}{\partial r} \right) + \frac{1}{\sin \theta} \frac{\partial}{\partial \theta} \left(\sin \theta \frac{\partial \Phi}{\partial \theta} \right) \\
& = -\Sigma [\lambda(|\Phi|^2 - F_0^2)\Phi],
\end{aligned} \tag{4.5}$$

where an overall factor of Σ is included to simplify the expression. Similar to the BTZ case, we extract the Coriolis and centrifugal terms by expanding in powers of Ω_K , to second order:

$$R_{\text{Coriolis}}^K = \frac{2\Omega_K}{g_{tt}} \frac{\partial^2 \Phi}{\partial t \partial \phi}, \quad R_{\text{centrifugal}}^K = \frac{\Omega_K^2}{g_{tt}} \frac{\partial^2 \Phi}{\partial \phi^2}. \tag{4.6}$$

These are formally the same as in the BTZ metric. We recover Eq. (3.12) when $g_{tt} \rightarrow -c^2$, in the limit $r \gg r_s$ and $c \rightarrow \infty$. These quantities vanish for the Schwarzschild metric, for which $\Omega_K = 0$. In our calculations, we use the potential approximation to the unperturbed Schwarzschild metric [25].

In the slow-rotation regime, once we extract the terms R_{Coriolis} and $R_{\text{centrifugal}}$ from the Kerr black hole geometry to describe the rotation, the confining effect of the Kerr black hole on the superfluid is not very different from the case of the non-rotating Schwarzschild black hole. It is easy to see that Eq. (4.5) reduces to a NLKG in the Schwarzschild black hole background by setting $a = 0$; then, under the Newtonian approximation, an effective confining potential $\sim -\frac{M}{r}$ appears. Therefore, if M is large enough, rotating black holes can hold some amount of cosmic superfluid through their gravitational pull, and the effective potential plays the same role as the trapping potential in the rotating BEC context. We have not been able to solve Eq. (4.5) exactly. Nevertheless, a numerical computation based on the NLKG with a current-current interaction, which is used to simulate the local rotation terms R_{Coriolis} and $R_{\text{centrifugal}}$ of the Kerr black hole, has been performed in Ref. [21]. A solution of a three-dimensional vortex-ring lattice was found (see Fig. 4 in Ref. [21]), and it will be interesting to see whether such a solution exists for Eq. (4.5).

5 Conclusions and Outlooks

In this paper we have studied, from a mathematical point of view, the nonlinear Klein-Gordon equation in the (2+1)-dimensional BTZ black hole background and the (3+1)-dimensional Kerr black hole background. Physically speaking, we have studied the behavior of superfluids driven by rotating black holes (BTZ and Kerr) via the frame-dragging effect, and observed the formation of quantum vortex lattices by numerically solving the NLKG equation under tractable approximations (slow and constant rotation). In the case of slow rotation, we have identified Eq. (3.12), Eq. (3.16), and Eq. (4.6) as the relevant rotation terms for studying quantum vorticity in rotating backgrounds. The common form taken by these terms is explicitly shown and discussed. In addition, the vortex lattice in Fig. (1) has the same pattern as the lattices seen in rotating superfluid helium and rotating BECs [15, 16]. Although our results are derived in a restricted region of parameter space, they show how angular momentum in the metric can be transferred to the fields, resulting in vorticity and hence confirming superfluidity.

We do not speculate on whether quantum vorticity should be manifest in astrophysical observations. Our results confirm, only in principle, that rotating geometry induces

vorticity. Our purpose here was modest: to see if vorticity could be sourced by the angular momentum in the spacetime. As this has been confirmed, we will however remark that nucleation of quantum vortices in an astrophysical context could require significant frame-dragging, and this may be the case in the neighborhood of a black hole with ultra-high angular momentum. It has been speculated elsewhere that the so-called “non-thermal filaments” observed near the center of the Milky Way are due to quantized vorticity [7].

As a related issue, it is interesting to mention that as the angular momentum increases, one expects that vortex filaments would appear closer to the black-hole surface, forming a hydrodynamic boundary layer; this may be relevant to the problem of black-hole collapse. It is well known in hydrodynamics that such a boundary layer can accommodate any necessary boundary conditions for the surrounding fluid. One of the outstanding theoretical problems in general relativity is to extend the Oppenheimer-Snyder solution [26] of black-hole collapse in the Schwarzschild metric, which carries no angular momentum, to the Kerr metric. A main obstacle appears to be the lack of an appropriate generalization of the interior metric, which in the Oppenheimer-Snyder case is just the non-rotating closed Robertson-Walker metric. It is therefore tempting to contemplate that when a hydrodynamic boundary layer is present, the outside Kerr metric could be joined onto any interior metric.

The relativistic character of the curved-space NLKG equation is appropriate for addressing questions involving strong gravitational fields, such as those in the vicinity of the curved spacetime of a rotating black hole. Since quantum vorticity is synonymous with superfluids and is indicative of dynamical complex fields and non-linear pattern-forming systems, its presence in a gravitational context is particularly welcome. We hope that these investigations contribute to the productive and fascinating enterprise that has been the consideration of gravitational influence on quantum fields.

Acknowledgements

We thank Weizhu Bao and Yong Zhang of the National University of Singapore for helpful discussions and suggestions on numerical computations, and Yulong Guo for assistance in much of the coding. CX is supported by the research grant from the Institute of Advanced studies, Nanyang Technological University, Singapore and MRRG is funded in part by the Social Policy Grant at Nazarbayev University.

References

- [1] ATLAS collab., *Phys. Letters B* **716**, 1 (2012).
- [2] CMS collab., *Phys. Letters B* **716**, 30 (2012).
- [3] W. H. Zurek, *Nature* **317**, 505 (1985).
- [4] G. E. Volovik, *Czech. J. Phys.* **46**, 3048 (1996) [cond-mat/9607212].
- [5] G. E. Volovik, *Phys. Rept.* **351**, 195 (2001).
- [6] K. Huang, H. B. Low, and R. S. Tung, *Class. Quantum Grav.* **29**, 155014 (2012); arXiv:1106.5282.

- [7] K. Huang, H. B. Low, and R. S. Tung, *Int. J. Mod. Phys. A* **27**, 1250154 (2012); arXiv:1106.5283.
- [8] C. G. Böehmer and T. Harko, *J. Cosmol. Astropart. Phys.*, **06**, 025 (2007).
- [9] J.-W. Lee, *J. Korean Phys. Soc.* **54**, 2622 (2009).
- [10] V. Popov, *Phys. Lett. B* **686**, 211 (2010).
- [11] B. Kain and H.Y. Ling, *Phys. Rev. D* **82**, 064042 (2010).
- [12] A. Suarez, V.H. Robles, and T. Matos, *Astrophys. Space Sci. Proc.* **38**, 107 (2014) [arXiv:1302.0903 [astro-ph.CO]].
- [13] K. Huang, C. Xiong, and X. Zhao, *Int. J. Mod. Phys. A* **29**, 1450014 (2014); arXiv:1304.1595.
- [14] K. Huang, *Int. J. Mod. Phys. A* **28**, 1330049 (2013); arXiv:1309.5707.
- [15] R. J. Donnelly, *Quantized vortices in helium II* (Cambridge University Press, 1991).
- [16] For a review, see e.g. A. L. Fetter, *Rev. Mod. Phys.* **81**, 647 (2009).
- [17] R. P. Kerr, *Phys. Rev. Lett.* **11**, 237 (1963).
- [18] M. Bañados, C. Teitelboim, and J. Zanelli, *Phys. Rev. Lett.* **69**, 1849 (1992).
- [19] J. Lense and H. Thirring, *Phys. Zeits.* **19**, 156 (1918).
- [20] C. Everitt *et. al.*, *Phys. Rev. Lett.*, 106, 221101 (2011); arXiv:1105.3456.
- [21] C. Xiong, M. R. R. Good, Y. Guo, X. Liu and K. Huang, *Phys. Rev.* **12**, D90, 125019 (2014). arXiv:1408.0779 [hep-th].
- [22] M. Kenmoku, M. Kuwata, K. Shigemoto, *Class.Quant.Grav.* **25**:145016, (2008).
- [23] M. R. R. Good and Y. C. Ong, *Phys. Rev. D* **91**, no. 4, 044031 (2015) [arXiv:1412.5432 [gr-qc]].
- [24] R. M. Wald, *General Relativity* (University of Chicago Press, 1984), p.319.
- [25] C. W. Misner, K. S. Thorne, and J. A. Wheeler, *Gravitation* (Freeman, San Francisco, 1973), Chap.25, p.636.
- [26] J. R. Oppenheimer and H. Snyder, *Phys. Rev.* **56**, 455 (1939).