

## ASTROPROTECTIVE POTENTIAL OF EXTRACT FROM LIMONIUM GMELINII FOR STROKE THERAPY

Shayakhmetov Y., Tsoy A., Zhussupova G., Askarova Sh.

*National Laboratory Astana, Nazarbayev University (Astana, Kazakhstan)*  
[shaskarova@nu.edu.kz](mailto:shaskarova@nu.edu.kz)

**Background:** It's known that proinflammatory cytokines, such as TNF- $\alpha$ , produced by injured neurons and infiltrated leukocytes, involved in every aspect of stroke. Cytokines promote activation of astroglia, induce oxidative stress thereby exacerbating brain tissue damage. There is a number of data indicating that polyphenols can provide protection against neurodegenerative changes. It has been reported previously, that reach with polyphenols extract of *Limonium gmelinii* exerts a wide range of therapeutic actions. Here, we aimed to study antioxidant and anti-inflammatory potential of *L. gmelinii* under exposure of TNF- $\alpha$  in vitro.

**Materials and methods:** In this study we used human primary astrocytes as following: control; cells incubated with TNF- $\alpha$  (0.1 ng/ml for 4 hr); cells pretreated with plant extract of *L. gmelinii* (30 ug/ml for 18 hr) followed by TNF- $\alpha$  exposure; cells incubated with plant extract only. To quantify the NADPH subunits colocalization, Erk 1/2 activation and ROS generation in astrocytes we applied quantitative fluorescence microscopy, confocal microscopy and western blotting.

**Results:** We observed that TNF- $\alpha$  promoted NADPH oxidase activation and consequent ROS generation and Erk 1/2 phosphorylation. In contrast, incubation of astrocytes with *L. gmelinii* extract suppressed TNF- $\alpha$  -induced colocalization between the p47-phox and gp91-phox subunits of NADPH oxidase, ROS production and phosphorylation of ERK1/2. At the same time, plant extract alone did not affect ROS formation and activity of the enzymes.

**Conclusion:** Our results demonstrate that extract from *L. gmelinii* possesses antioxidant and anti-inflammatory properties in vitro. Therefore, further studies are required to assess the effectiveness of the *L. gmelinii* extract in vivo.