

Original Article

Existence of self-similar solutions of the two-dimensional Navier–Stokes equation for non-Newtonian fluidsDONGMING WEI^a, SAMER AL-ASHHAB^{b,*}^a Department of Mathematics, Nazarbayev University, 53 Kabanbay Batyr Ave, Astana, 010000, Kazakhstan^b Department of Mathematics and Statistics, Al Imam Mohammad Ibn Saud Islamic University (IMSIU), P.O. Box 90950, Riyadh 11623, Saudi Arabia

Received 23 October 2018; revised 12 March 2019; accepted 6 April 2019

Available online xxxx

Abstract. The reduced problem of the Navier–Stokes and the continuity equations, in two-dimensional Cartesian coordinates with Eulerian description, for incompressible non-Newtonian fluids, is considered. The Ladyzhenskaya model, with a non-linear velocity dependent stress tensor is adopted, and leads to the governing equation of interest. The reduction is based on a self-similar transformation as demonstrated in existing literature, for two spatial variables and one time variable, resulting in an ODE defined on a semi-infinite domain. In our search for classical solutions, existence and uniqueness will be determined depending on the signs of two parameters with physical interpretation in the equation. Illustrations are included to highlight some of the main results.

Keywords: Non-linear boundary value problem; Singular; Self-similar transformation; Existence; Uniqueness

Mathematics Subject Classification: 34B40; 76A05

1. INTRODUCTION

The study of non-Newtonian fluids, both mathematically and physically, has gained much importance during the last few decades due to their many applications in industry

* Corresponding author.

E-mail addresses: dongming.wei@nu.edu.kz (D. Wei), ssashhab@imamu.edu.sa (S. Al-Ashhab).

Peer review under responsibility of King Saud University.

**Production and hosting by Elsevier**

<https://doi.org/10.1016/j.ajmsc.2019.04.001>

1319-5166 © 2019 The Authors. Production and Hosting by Elsevier B.V. on behalf of King Saud University. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

and in describing physical phenomena. The basic physical theory, and its mathematical formulation can be found in [1,8,18]. Many researchers studied non-Newtonian fluids from a numerical or computational point of view, in some instances accompanied with certain techniques or transformations to elucidate investigating the problem [6,9]. Other studies involved existence and uniqueness of solutions to problems involving non-Newtonian fluids [10,11,20,21]. Many times, it is found that solutions for Newtonian and non-Newtonian flows are not unique [7,13,15,17]. In some instances or special cases, exact solutions were established, see for example [12]. Our interest in this paper is in a Ladyzhenskaya type non-Newtonian fluid [16], where self-similar transformations of the Navier–Stokes equations, for non-Newtonian incompressible fluids, lead to an ODE with dependence on one similarity variable. Navier–Stokes equations in two dimensions, for incompressible non-Newtonian fluids, consist of a system of PDEs with two spatial variables, and a time variable. However, a two-dimensional generalization of the well-known self-similar Ansatz reduces the PDE system into an ODE. This resulting ODE was used for example in [4], to study the compressible Newtonian Navier–Stokes equations. Symmetry reductions analysis can also be applied to obtain some solutions, as was done in [14], and as was done for three dimensions in [19].

Recently in [3], the authors considered a self-similar transformation to obtain analytic solutions of the two-dimensional Navier–Stokes equations, with Eulerian description, for a non-Newtonian fluid. However, it remains to investigate existence and uniqueness of solutions for that particular reduced Navier–Stokes equation, with suitable boundary conditions. A similar problem was studied in [5], but where the parameters were tied together via certain relations, and where the authors used a different approach to investigate the problem.

We shall discuss existence (or non-existence) and uniqueness of solutions for the resulting Navier–Stokes reduced problem. In Section 2, we introduce the problem with a brief derivation including the main ideas leading to the governing equation of interest. The main results are then derived in Section 3, where we discuss separate cases depending on the sign of two parameters: the flow behavior index (mathematically an exponent r) and the leading coefficient k in the governing equation.

2. THE PROBLEM

Consider the Ladyzhenskaya model of non-Newtonian fluid dynamics, with the following formulation (c.f. [16]):

$$\rho \frac{\partial \mathbf{u}_i}{\partial t} + \rho \mathbf{u}_j \frac{\partial \mathbf{u}_i}{\partial x_j} = -\frac{\partial p}{\partial \mathbf{x}_i} + \frac{\partial \Gamma_{ij}}{\partial x_j} + \rho \mathbf{F}_i \quad (1)$$

$$\frac{\partial \mathbf{u}_j}{\partial x_j} = 0 \quad (2)$$

where the Einstein summation convention is assumed on the j index. The parameters ρ , \mathbf{u} , p and \mathbf{F} represent the density, the two dimensional velocity field, the pressure, and the external force, respectively. On the other hand, observe that Γ_{ij} is defined via:

$$\Gamma_{ij} = (\mu_0 + \mu_1 |E(\nabla \mathbf{u})|^r) E_{ij}(\nabla \mathbf{u}) \quad (3)$$

where μ_0 , μ_1 and r represent the dynamical viscosity, the consistency index, and the flow behavior index, respectively, and where

$$E_{ij}(\nabla \mathbf{u}) = \frac{1}{2} \left(\frac{\partial \mathbf{u}_i}{\partial x_j} + \frac{\partial \mathbf{u}_j}{\partial x_i} \right) \quad (4)$$

is the Newtonian linear stress tensor. Observe that \mathbf{x} represents the two dimensional Cartesian coordinates, say $\mathbf{x} = (x, y)$. Now, setting the external force to zero $\mathbf{F} = \mathbf{0}$, observing that in two dimensions:

$$|E| = (u_x^2 + v_y^2 + \frac{1}{2}(u_y^2 + v_x^2))^{1/2},$$

(where u and v are the components of \mathbf{u}) and letting:

$$L = \mu_0 + \mu_1|E|^r,$$

simplifies the formulation, using compact notation, to the following equations:

$$u_x + v_y = 0, \tag{5}$$

$$u_t + uu_x + vv_y = -\frac{p_x}{\rho} + L_x u_x + Lu_{xx} + \frac{L_y}{2}(u_y + v_x) + \frac{L}{2}(u_{yy} + v_{xy}), \tag{6}$$

$$v_t + uv_x + vv_y = -\frac{p_y}{\rho} + L_y v_y + Lv_{yy} + \frac{L_x}{2}(u_y + v_x) + \frac{L}{2}(v_{xx} + u_{xy}). \tag{7}$$

The following transformation (8) (self-similar Ansatz, c.f. [3]) leads to solutions of physical interest, and shall further simplify the problem consisting of the 3×3 PDE system (5)–(7) given above. Namely, this transformation is given by:

$$u = t^{-\alpha} f(\eta), \quad v = t^{-\beta} g(\eta), \quad p = t^{-\gamma} h(\eta), \quad \eta = t^{-\delta}(x + y) \tag{8}$$

where η is called a similarity variable. The functions f, g , and h are referred to as shape functions. We shall consider $\mu_0 = 0, \mu_1 \neq 0$, and we note that the details of the entire derivation and simplification process can be found in the references, c.f. [2,3] and the references therein. We choose to skip those details since our main interest is in the resulting ODE for f below. However, we do point out that through the simplification process, the shape functions are assumed to have interrelations relating them to one another, while the following relations are obtained for the above exponents:

$$\alpha = \beta = (1 + r)/2, \quad \delta = (1 - r)/2, \quad \gamma = r + 1. \tag{9}$$

Solutions of physical relevance and interest will require all exponents in (9) to be positive, from which we must have: $-1 < r < 1$. It is noted that in similar power-law problems, a power-law index n is used and is related to r mathematically via $r = n - 1$. In this respect, $-1 < r < 0$ corresponds to pseudo-plastic or shear-thinning fluid, while $0 < r < 1$ corresponds to a shear-thickening fluid. (Since $r > 1$ has been eliminated, the fluid of interest here maybe considered as a restricted Ostwald–de Waele-type fluid.) The following ODE is the reduced and simplified equation that is of our interest, and it is the following reduced Navier–Stokes equation:

$$2^{r+1}(1 + r)\mu_1 f''|f'|^{r-1} f' + (1 - r)\eta f' + (1 + r)f = 0. \tag{10}$$

Observe that this ODE is for f , while g and h are related to f via certain relations as can be found in the references. Due to the conditions we shall consider, see (12), we shall suppose $f' \leq 0$. (Observe that if f' reaches zero at some point, say $f'(\eta_0) = 0$, then the equation may become inconsistent in case $f(\eta_0) \neq 0$ for $r > 0$, or it may become undefined if $r < 0$.) By further assuming

$$k = 2^{r+1}(1 + r)\mu_1,$$

we obtain the equivalent equation (11). Before proceeding with the analysis, however, observe that if $f'(\eta_0) = 0$ while $f(\eta_0) \neq 0$, for some $\eta_0 > 0$, then Eq. (10) becomes inconsistent for positive r . The solution assumes a point of termination at such instances. Solutions also assume a terminal point for negative values of r when $f'(\eta_0) = 0$ as the first term in the ODE becomes undefined. It is noted that practical values of $k > 0$ were listed in [3], while $k < 0$ can be found in the similar Rayleigh problem. So, now, consider:

$$-kf''(-f')^r + (1-r)\eta f' + (1+r)f = 0 \tag{11}$$

We shall make a few observations regarding (11). First, notice that if $r = 0$ then we have the equation $-kf'' + (\eta f)' = 0$ which leads to a solution: $-kf' + \eta f = c$ and therefore $f(\eta) = f(0)e^{\eta^2/2k} + f'(0)e^{\eta^2/2k} \int_0^\eta e^{-u^2/2k} du$. This solution approaches zero for $k < 0$ as $\eta \rightarrow \infty$, and consequently it is an explicit illustration of the existence of a solution when $r = 0, k < 0$, which satisfies (12).

Additionally, observe that it is not possible to have $f \rightarrow c \neq 0$ as $\eta \rightarrow \infty$, for some constant $c \neq 0$, unless f reaches c at some finite η . To establish this, let $g(\eta) = f(\eta) - c$ so that $f(\eta) = c + g(\eta)$, then we must have $g(\eta) \rightarrow 0$ as $\eta \rightarrow \infty$, and therefore $-kg''(-g')^r + (1-r)\eta g' = -(1+r)(c + g)$, which upon integration would imply that:

$$\frac{k(-g'(\eta))^{r+1}}{r+1} = -(1+r)c\eta - (1-r)\eta g(\eta) - 2r \int_0^\eta g(u)du + K,$$

where $K = \frac{k(-g'(0))^{r+1}}{r+1}$ is a constant. Now, since $r > -1$ and the first term on the right-hand side would make that side of the equation diverge and become unbounded as $\eta \rightarrow \infty$, this would in turn imply that the equation does not balance, or otherwise $g'(\eta)$ has to take on infinite values as $\eta \rightarrow \infty$, which is a contradiction. It is very important to emphasize here that it will be shown that solutions do exist where f reaches $c \neq 0$ at a terminal point in finite η : $f(\eta_0) = c \neq 0, f'(\eta_0) = 0$ for some $\eta_0 > 0$, as is also shown in numerical illustrations in [3] for $r < 0$. The boundary conditions for an equation such as (11) are typically given at 0 and at ∞ . The boundary conditions of interest to us take the form:

$$f(0) = a, \quad f(\infty) = 0 \tag{12}$$

where $a > 0$.

3. EXISTENCE OF SOLUTIONS

To establish *existence of solutions*, a shooting method is utilized where the condition at infinity is replaced by an initial condition $f'(0)$: we shall first show that Eq. (11) subject to $f(0) = a$ (the first of the two conditions in (12)) has solutions for which $f'(\eta_0) = 0$ at some finite $\eta_0 < \infty$ and where $f(\eta_0) = b > 0$ (such solutions terminate at η_0 as discussed above) for some appropriate choice of $f'(0)$. We shall also show that it has solutions that extend to infinite η while crossing the horizontal axis at some point.

Observe that subtracting $2rf$ from both sides of Eq. (11) yields the following: $-kf''(-f')^r + (1-r)\eta f' + (1-r)f = -2rf$, where now observe that the left-hand side is an exact derivative. Now integrating from 0 to η and using a dummy variable of integration, say t , we obtain

$$(-f'(\eta))^{r+1} = (-f'(0))^{r+1} - \frac{(r+1)}{k} \left((1-r)\eta f(\eta) + 2r \int_0^\eta f(t)dt \right). \tag{13}$$

To begin with, let us consider the case $r > 0, k > 0$:

Theorem 1. *There exists a unique solution to (11) subject to (12) for $r > 0, k > 0$, and where $f(\eta) > 0$ for all $\eta > 0$.*

Proof. To begin with, we show that for some appropriate choice of the initial condition $f'(0) < 0$ one obtains a solution that terminates at some finite η_0 where $f'(\eta_0) = 0, f(\eta_0) > 0$. Observe that (11) implies that $f''(0) > 0$. We further assume $f'' > 0$ on the entire interval $(0, \eta_0)$ which will be verified at the end of the proof, and with $f'' > 0$ we must have:

$$(-f'(\eta))^{r+1} < (-f'(0))^{r+1} - \frac{(r+1)}{k} \left(0 + 2r \int_0^\eta (f(0) + f'(0)t) dt \right),$$

and therefore

$$(-f'(\eta))^{r+1} < (-f'(0))^{r+1} - \frac{2r(r+1)}{k} (f(0)\eta + f'(0)\eta^2/2).$$

Taking $(-f'(0))^{r+1} < \frac{r(r+1)}{k} f(0)$ and $|f'(0)| < f(0)$ (whichever yields a smaller $|f'(0)|$, recall that $f'(0)$ is negative) would in fact show that for $\eta = 1$ we have $(-f'(1))^{r+1} < 0$, but by assumption this last quantity should be non-negative (due to $f' < 0$). This contradiction shows that $f' = 0$ at some finite $\eta_0 < 1$. Finally one checks that with the additional condition $|f'(0)| < \frac{(r+1)}{2} f(0)$ we have $f'' > 0$ and $f > 0$ for all $\eta < 1$, so that the above arguments hold (note that this strong condition for $|f'(0)|$ establishes our point here, but it might be relaxed significantly once a particular solution is determined).

On the other hand, it can be shown that for large enough $|f'(0)|$ we obtain a solution for which $f'(\eta) < 0$ for all $\eta > 0$, and where $f(\eta) < 0$ for all $\eta > \eta_0$, for some $\eta_0 > 0$ (i.e. a solution that crosses the η -axis). Now observe that for $f' < 0$ it follows from Eq. (11) that $-kf''(-f')^r = -(1-r)f' - (1+r)f > -(1+r)f$, which can be integrated to obtain

$$(-f'(\eta))^{r+1} > (-f'(0))^{r+1} - \frac{(r+1)}{k} \int_0^\eta f(t) dt, \tag{14}$$

from which we have

$$(-f'(\eta))^{r+1} > (-f'(0))^{r+1} - \frac{(r+1)}{k} f(0)\eta; \tag{15}$$

by choosing $f'(0)$ to be large enough in absolute value such that

$$(-f'(0))^{r+1} > (f(0))^{r+1} + \frac{(r+1)}{k} f(0) \tag{16}$$

then it is guaranteed from (15) and (16) that $(-f'(\eta))^{r+1} > (f(0))^{r+1}$ for all $0 < \eta < 1$, and therefore $f'(\eta) < -f(0) < 0$ for all $0 < \eta < 1$, which in turn guarantees the existence of some $\eta_0 < 1$ such that $f(\eta_0) = f(0) + \int_0^{\eta_0} f'(t) dt = 0$. Once we have $f(\eta_0) = 0$ with $f'(\eta_0) < 0$, then Eq. (11) will show that this solution will satisfy: $f(\eta) < 0, f'(\eta) < 0$ for all $\eta > \eta_0$. (We note that the same argument can be used for $-1 < r < 0$ since the exponent $r + 1$ is positive for this range of r , as will be needed for later proofs.)

Now to show existence of solutions: given the above results, suppose that y_1 is a solution that terminates at some finite η_1 where $y_1'(\eta_1) = 0$ and $y_1(\eta_1) = \epsilon > 0$. One can find another solution that terminates at $y_2(\eta_2) = \epsilon/2$ for some η_2 , i.e., $y_2(\eta_2) = \epsilon/2, y_2'(\eta_2) = 0$. It is not difficult to prove this last mathematical statement, following similar analysis as above, coupled with the continuity with respect to initial conditions (on the interval $(0, \eta_1)$). We, however, leave out some of the obvious details.

In fact, a general assumption that there is a *minimum* value for a solution $f > 0$ where f' reaches zero so the solution terminates (at say η_1 , i.e. $f(\eta_1) = \epsilon_{\min} > 0, f'(\eta_1) = 0$, and where no solution with smaller f -values will terminate), leads to a contradiction for the case $r > 0, k > 0$. Since then, one can still take a *slightly* larger $|f'(0)|$ so that $f(\eta_1)$ decreases very slightly, while the new $|f'(\eta_1)|$ is very small so that $f(\eta)$ will still have to decrease for $\eta > \eta_1$. But on the other hand, $f''(\eta)$ would be *large enough* for $\eta > \eta_1$, and will approach infinity fast since $r > 0$, see (11). The new solution will then terminate with a smaller $f > 0$ at say $\eta_2 > \eta_1$.

We still need to prove that there exists a solution that will not reach $f = 0$ at finite η , i.e., we need to show that $f \rightarrow 0$ with $f > 0$ for all $\eta > 0$.

So now with $y_2(\eta_2) = \epsilon/2$ as above, observe that if we let $\delta_2 = (-y_2'(0))^{r+1}$, where $y_2(0)$ is the initial condition corresponding to the solution y_2 , which is extended to, and terminates at η_2 , then Eq. (13) yields the following: $\delta_2 = \frac{(r+1)}{k}((1-r)\eta_2(\frac{\epsilon}{2}) + 2r \int_0^{\eta_2} y_2(t)dt)$ since $(-y_2'(\eta_2))^{r+1} = 0$. Similarly $\delta_1 = \frac{(r+1)}{k}((1-r)\eta_1\epsilon + 2r \int_0^{\eta_1} y_1(t)dt)$, where $\delta_1 = (-y_1'(0))^{r+1}$, and y_1 is the solution extending to η_1 with $y_1(\eta_1) = \epsilon, y_1'(\eta_1) = 0$. Therefore

$$\delta_2 - \delta_1 = \frac{(r+1)}{k} \left(\epsilon(1-r)(\frac{\eta_2}{2} - \eta_1) + 2r \int_0^{\eta_1} (y_2(t) - y_1(t))dt + 2r \int_{\eta_1}^{\eta_2} y_2(t)dt \right).$$

Observe that the last two terms in parentheses on the right-hand side of the equation above satisfy:

$$2r \int_0^{\eta_1} (y_2(t) - y_1(t))dt + 2r \int_{\eta_1}^{\eta_2} y_2(t)dt < \frac{3\epsilon}{2}(\eta_2 - \eta_1),$$

since the first integral is negative, and the second integral is smaller than the trapezoidal area under the line extending between (η_1, ϵ) and $(\eta_2, \epsilon/2)$. This area is equal to $\frac{3\epsilon}{4}(\eta_2 - \eta_1)$, and after multiplying this area by $2r$ and recalling that $0 < r < 1$, the desired result is obtained. Now, note that $\delta_2 - \delta_1 > 0$ so we can deduce that $\epsilon(1-r)(\frac{\eta_2}{2} - \eta_1) + \frac{3\epsilon}{2}(\eta_2 - \eta_1) > 0$, and therefore $\frac{\eta_2}{\eta_1} > \frac{5-2r}{4-r} = K > 1$, for $0 < r < 1$. In this manner, it can be shown that the solution can be extended to $\eta = \infty$ since we can go step by step to $y = \epsilon/2^n, n = 1, 2, 3, \dots$, and reach $\eta > K^n \eta_1$, where $K = \frac{5-2r}{4-r} > 1$ as given above.

To verify that f'' stays negative for the new solution y_2 one can check that $f''' = \frac{-f'(\eta f''(1-r)^2 + 2f') + r(1+r)ff''}{k(-f')^{r+1}}$. So, on the one hand, if $y_2(\eta_1)$ goes significantly below ϵ , with $y_2'(\eta_1)$ relatively small in absolute value so that $y_2''(\eta_1)$ is large, and f'' approaches infinity quickly, then it is obvious that f'' stays positive (from (11)). On the other hand, if $y_2(\eta_1)$ goes slightly below ϵ , say to ϵ_0 , with $y_2'(\eta_1)$ becoming relatively large in absolute value, then keep δ_2 small, or close enough to δ_1 , so that $y_2'(\eta_1) = \frac{-\epsilon_0(1+r)}{\eta_1(1-r)} + \epsilon'$ for some very small ϵ' that will yield $y_2''(\eta_1) = \frac{-2y_2'(\eta_1)}{\eta_1(1-r)^2}$ from (11). Observe now that the above expression for f''' is positive at η_1 (with both terms in the numerator being positive) and will stay positive with f'' increasing, and f' increasing (becoming closer to zero). The fact that now $y_2''(\eta_1)$ is relatively *very small* and using the above expression for f''' , shows that by the point where we get to a terminal point with $y_2' = 0$ and y_2'' becoming unbounded, it must be that y_2 is significantly smaller than ϵ , and where we leave out some of the details. The process can be repeated to eventually get to a solution where $y_2(\eta_2) = \epsilon/2$ and where $y_2' > 0$ is




Fig. 1. A typical solution to the Navier–Stokes equation (11) with $r > 0, k > 0$.

guaranteed on the maximal interval of continuation for y_2 . Observe that this also reinforces our earlier discussion on the existence of y_2 reaching $\epsilon/2$ and terminating.

To establish uniqueness, suppose that $f(\eta)$ is a solution that satisfies (11) subject to (12). Define

$$F(\eta) = \frac{(r + 1)}{k} \left((1 - r)\eta f(\eta) + 2r \int_0^\eta f(t)dt \right),$$

and note that $F(\eta)$ is an increasing function such that in the limit we have: $F(\eta) \rightarrow (-f'(0))^{r+1}$ as $\eta \rightarrow \infty$, and where $f'(0)$ is the initial condition corresponding to the given solution f . Suppose that $g(\eta)$ is another solution with $g'(0) \neq f'(0)$, say $(-g'(0))^{r+1} = (-f'(0))^{r+1} + \epsilon$ with $\epsilon \neq 0$. Take $\epsilon > 0$: the solution g will then satisfy $g(\eta) < f(\eta)$, for all $\eta > 0$, so that:

$$G(\eta) = \frac{(r + 1)}{k} \left((1 - r)\eta g(\eta) + 2r \int_0^\eta g(t)dt \right) \leq F(\eta), \tag{17}$$

and where $G(\eta) \rightarrow (-g'(0))^{r+1} = (-f'(0))^{r+1} + \epsilon$ as $\eta \rightarrow \infty$, which follows from our assumption that g is another solution that satisfies (12). But then we would have $G(\infty) > (-f'(0))^{r+1} = F(\infty)$, and this last inequality requires $G(\eta) > F(\eta)$ for large η , which is a contradiction (it contradicts (17)). This completes the proof.

Fig. 1 shows a typical solution to the Navier–Stokes equation (11) illustrating the above result. Another result can readily be obtained here for $r > 0, k < 0$:

Proposition 2. *There exists no solution to (11) subject to (12) for $r > 0, k < 0$ and where $f(\eta) \geq 0$ for all $\eta > 0$.*

Proof. Under the hypotheses of the preceding theorem where $f(\eta) > 0$ for all η , Eq. (13) will show that $(-f'(\eta))^{r+1} > (-f'(0))^{r+1} > 0$. This implies that it is not possible to have $f \rightarrow 0$ as $\eta \rightarrow \infty$. Nor is it possible to have a solution that reaches zero equilibrium at finite η : $f'(\eta) = 0$ when $f(\eta) = 0$, for the same reason.




Fig. 2. A typical solution to Eq. (11) with $r > 0, k < 0$. It crosses the axis.

In fact, solutions where $r > 0, k < 0$, will cross the axis, and will eventually terminate at some point where $f'(\eta_0) = 0, f(\eta_0) < 0$, for some finite η_0 . This can be illustrated with the aid of numerical integrators. (See Fig. 2.)

3.1. The case $r < 0, k > 0$

As for the case where $r < 0, k > 0$, we begin by showing that a solution exists where $f'(\eta_0) = 0$ at some finite $\eta_0 > 0$: observe that with $f' < 0, f'' > 0$ we have $f(\eta) > f(0) + f'(0)\eta$, so that Eq. (13) yields:

$$\begin{aligned} (-f'(\eta))^{r+1} &< (-f'(0))^{r+1} - \frac{(r+1)}{k} ((1-r)(f(0) + f'(0)\eta)\eta + 2rf(0)\eta) \\ &< (-f'(0))^{r+1} - \frac{(r+1)}{k} ((1+r)f(0)\eta + (1-r)f'(0)\eta^2). \end{aligned}$$

Choose $f'(0)$ small enough in absolute value so that:

$$f(0) > \frac{k}{(1+r)^2} (-f'(0))^{r+1} - \frac{(1-r)}{(1+r)} f'(0).$$

This choice will show that a solution exists such that for some $\eta_0 < 1$, we have $f'(\eta_0) = 0$, and the solution terminates. It can readily be verified that $f' < 0, f'' > 0$, within the interval of the given solution, so that the above arguments stay valid.

On the other hand, there exists a solution which crosses the axis at some finite η . This can be established using the same arguments in the proof of the preceding theorem, as was stated earlier. However, observe that since $k > 0$ and $f' < 0$, we must have

$$(1-r)\eta f' + (1+r)f > 0 \tag{18}$$

in order to avoid any inflection point (with $f > 0$, and since the solution will cross the axis once it has an inflection point, as the curvature will continue to be negative once it is negative). Observe, now, that inequality (18) implies $\frac{f'}{f} > -\frac{(1+r)}{(1-r)\eta}$, and therefore




Fig. 3. A set of solutions to Eq. (11) with $r < 0, k > 0$. They do not satisfy (12): There is a minimum for f where f' reaches zero and (11) becomes undefined (a terminal point), beyond which solutions change curvature with $f'(\eta) < 0$ on the entire solution domain.

$f > c\eta^{-\frac{(1+r)}{(1-r)}}$, where c is a constant, and $-\frac{(1+r)}{(1-r)} < 0$ for $-1 < r < 1$. Now, if $f = \eta^p$ where $p > -\frac{(1+r)}{(1-r)}$, then the above inequality for f holds, but inequality (13) will have a divergent term on the right-hand side, and therefore f' will reach zero in finite time say η_1 , with $f(\eta_1) > 0$, so that conditions (12) will not be satisfied. On the other hand, if we let $f(\eta) = c\eta^{-\frac{(1+r)}{(1-r)}} + g(\eta)$, with $0 < g(\eta) < \eta^q$ (of order q less than $p = -\frac{(1+r)}{(1-r)}$, q is real and $q < p$) then the above inequality still holds, but again a contradiction occurs upon substituting into (11), where we are led again to obtaining an inflection point. Therefore:

Theorem 3. *There exists no solution to (11) subject to (12) if $r < 0$, and $k > 0$.*

The dynamics here is the following: Solutions exist where f' reaches zero at some $\eta_0 > 0$, and $f(\eta) = b$ for all $\eta_0 < \eta < \infty$, for some large enough $b > 0$. However, there exists a certain value for $b > 0$ where further reduction of the initial condition $f'(0)$ (increase in absolute value of the gradient) shall yield a solution that crosses the horizontal axis ($f'(\eta)$ does not reach zero but rather stays negative). This happens since the decay of solutions (changes in f and f') becomes extremely slow with f'' proportional to $((1-r)\eta f' + (1+r)f)(f')^{-r}$ (namely observe the factor $(f')^{-r}$ with $f' \approx 0$ and where now $r < 0$), allowing the non-autonomous term $(1-r)\eta f'$ with the presence of η , to exceed the last term $(1+r)f$, of the governing equation (11). This leads to a change in curvature, and therefore solutions will cross the axis, and will not satisfy $f(\infty) = 0$ from (12). This is verified by numerical integrators, and is illustrated in Fig. 3: In particular the two upper curves reach a point where (11) is undefined with $f' = 0$. Such solutions reach a terminal point, that they cannot be extended beyond. The solution in the bottom illustrates that there is a minimum for f with those terminal points, after which solutions change curvature, and eventually will cross the axis.




Fig. 4. A typical solution to Eq. (11) with $r < 0, k < 0$. It reaches zero equilibrium at finite η (≈ 30 in this particular figure).

3.2. The case $r < 0, k < 0$

Unlike some of the previous cases, observe that in this case the governing equation (11) implies that $f''(0) < 0$. In fact, the curvature stays negative for some interval say $(0, \eta_0)$, until $f(\eta)$ drops in value while $f'(\eta)$ becomes more negative (see (11)). Then $f''(\eta)$ becomes positive, and it can readily be established that $f''(\eta)$ stays positive, on the infinite interval, if $|f'(0)|$ is large enough. Additionally, if the solution crosses the horizontal axis then $f''(\eta)$ will continue to be positive in this case of $k < 0$, and in fact if the solution does cross the axis it will eventually terminate with $f' = 0$: once the solution attains a negative value, say f_0 , then we have $f''(-f')^r > (1+r)f_0/k$, so that $-(-f'(\eta))^{r+1} \geq ((1+r)^2 f_0/k)(\eta - \eta_0) - (-f'(\eta_0))^{r+1}$, which implies that $f'(\eta)$ will reach zero at finite η . With the existence of solutions that cross the axis and then reach $f' = 0$, as stated by the remarks given above, another result is needed:

Lemma 4. *Two different solutions of (11) with the same initial $f(0)$, but two different initial gradients $f'_1(0) \neq f'_2(0)$, do not intersect for any $\eta > 0$. Furthermore, if $f'_2(0) < f'_1(0)$ with $f_2(0) \leq f_1(0)$, then $f_2(\eta) < f_1(\eta)$ for all $\eta > 0$.*

Proof. Given a solution with say $f'_1(0)$, take another solution with $f'_2(0) < f'_1(0)$, and where $f_2(0) = f_1(0)$. The two solutions will be different in, at least a small interval say $(0, \eta_0)$, and $f_2 < f_1$ on that interval. If the two solutions intersect, then $\eta f(\eta)$ would be the same for f_1 and f_2 at the point of intersection, and therefore the right-hand side of (13) would be larger for the solution f_2 . This, in turn, implies that $f_2(\eta)$ is larger than $f_1(\eta)$ in absolute value, so that $f'_2(\eta) < f'_1(\eta)$ at the point of intersection, and now this is a contradiction (which in fact can also be illustrated geometrically, as well as analytically).

Now, using the continuity with respect to initial conditions, it can be concluded that the solution f_2 with the larger initial absolute gradient $|f'_2(0)| > |f'_1(0)|$ will always

have a larger $|f_2'(\eta)|$, at all $\eta > 0$ where $f_1'(\eta) < 0$ (i.e. avoiding a situation where $f'(\eta) = 0$). Otherwise, at an η where $f_2'(\eta) = f_1'(\eta)$, let us say that $\epsilon > 0$ represents the difference between the two solutions: $f_2(\eta) = f_1(\eta) - \epsilon$. Then, observe that we would have $f_2''(\eta) > f_1''(\eta)$, where $f_2''(\eta)$ is larger precisely by the amount $\epsilon(1+r)(-f')^{-r}/k$ (see (11)). Now we can take ϵ small enough so that the two solutions would intersect at some point, say at $\eta + \Delta\eta$ (an argument here can be made, for example, using a Taylor series expansion). This contradicts the first result in the lemma, proven above. Now, note that the possibility $f_2'(\eta) > f_1'(\eta)$ would imply that $f_2'(\eta_0) = f_1'(\eta_0)$ at some $0 < \eta_0 < \eta$, since $f_2'(0) < f_1'(0)$. Therefore, the obtained contradiction would still eliminate this last possibility. This result can be generalized using similar arguments for $f_2(0) < f_1(0)$.

With solutions that reach $f' = 0$, $f = \text{constant} < 0$, and the above lemma, we may “construct” a solution that reaches zero equilibrium ($f = 0$) at finite η : given a solution that reaches equilibrium at a constant $f = c < 0$, take another solution with a smaller $|f'(0)|$ so that it reaches a terminal point $f = d > c$, at a *smaller* value of η (with $f'(\eta) = 0$). (This is a consequence of the preceding lemma.) Proceed in this fashion to find a solution that reaches zero at finite η (See Fig. 4.). Another way to view this is the following: we have solutions that cross the horizontal axis at η_0 with a negative $f'(\eta_0)$, so that taking another solution with a smaller $|f'(0)|$ leads to a less negative $f'(\eta_0)$ at η_0 , and with $f(\eta_0) > 0$. If the change in $f'(0)$ is small enough, the new solution will then cross the axis, but at a larger η and with a smaller $|f'|$ (at the point of crossing). This process can be continued until the required solution is reached. So this solution is established here, mathematically, as a limiting case.

Remark. Observe that the two different views above involve the same set of solutions.

Theorem 5. *Solutions to (11) subject to (12) exist for $r < 0, k < 0$, and where $f(\eta) \geq 0$ for all $\eta > 0$.*

In fact, analysis of Eq. (13) suggests that other solutions may exist but where $f(\eta) > 0$ for all $\eta > 0$, and with possibly an infinite number of points where the solution changes curvature. In such a case, the quantity $\eta f(\eta)$ does not approach zero due to balancing positive and negative terms in (13), which cannot approach zero. Furthermore, it can be easily checked that any solution of (11), with $r < 0, k < 0, f(0) > 0$, and any choice of $f'(0) < 0$, will satisfy $f'(\eta) < 0$ for all $\eta > 0$ as long as $f(\eta) > 0$, and cannot approach an equilibrium $f = c > 0$.

4. CONCLUSIONS

We studied a reduced problem from the Navier–Stokes and the continuity equations in two-dimensional Cartesian coordinates, with Eulerian description, for incompressible non-Newtonian fluids. We have shown the existence of positive solutions to the reduced ODE, $f \geq 0, f' \leq 0$, and where $f(\infty) = 0$. Such solutions exist if $rk > 0$. Those solutions may not be unique if the flow behavior index $r < 0$. On the other hand, positive solutions do not exist if $rk < 0$. Additionally, a solution exists and has been explicitly expressed when $r = 0, k < 0$.

ACKNOWLEDGMENT

Dongming Wei is supported by the Kazakhstan Ministry of Education Grant # AP05134166.

REFERENCES

- [1] G. Astarita, G. Marrucci, *Principles of Non-Newtonian Fluid Mechanics*, McGraw-Hill, Malabar, 1974.
- [2] I.F. Barna, Self-similar solutions of three-dimensional Navier–Stokes equation, *Commun. Theor. Phys.* 56 (4) (2011) 745–750, <http://dx.doi.org/10.1088/0253-6102/56/4/25>.
- [3] I. Barna, G. Bognár, K. Hriczo, Self-similar analytic solution of the two-dimensional Navier–Stokes equation with a non-Newtonian type of viscosity, *Math. Model. Anal.* 21 (1) (2016) 83–94, <http://dx.doi.org/10.3846/13926292.2016.1136901>.
- [4] I.F. Barna, L. Matyas, Analytic solutions for the three-dimensional compressible Navier–Stokes equation, *Fluid Dyn. Res.* 46 (5) (2014) <http://dx.doi.org/10.1088/0169-5983/46/5/055508>.
- [5] N. Bedjaoui, M. Guedda, Z. Hammouch, Similarity Solutions of the Rayleigh problem for Ostwald-de Wael electrically conducting fluids, *Anal. Appl.* 9 (2) (2011) 135–159.
- [6] G. Bognár, Similarity solution of a boundary layer flow for non-Newtonian fluids, *Int. J. Nonlinear Sci. Numer. Simul.* 10 (2010) 1555–1566.
- [7] G. Bognár, On similarity solutions of boundary layer problems with upstream moving wall in non-Newtonian power-law fluids, *IMA J. Appl. Math.* 77 (2012) 546–562.
- [8] G. Bohme, Non-Newtonian fluid mechanics, in: *North-Holland Series in Applied Mathematics and Mechanics*, Amsterdam, 1987.
- [9] J.P. Denier, P. Dabrowski, On the boundary-layer equations for power-law fluids, *Proc. R. Soc. A* 460 (2004) 3143–3158.
- [10] W. Gao, J. Wang, Similarity solutions to the power-law generalized Newtonian fluid, *J. Comput. Appl. Math.* 222 (2008) 381–391.
- [11] M. Guedda, Z. Hammouch, Similarity flow solutions of a non-Newtonian power-law fluid flow, *Int. J. Nonlinear Sci.* 6 (3) (2008) 255–264.
- [12] M. Guedda, R. Kersner, Non-Newtonian pseudoplastic fluids: Analytical results and exact solutions, *Int. J. Non-Linear Mech.* 4 (7) (2011) 949–957.
- [13] Z. Hammouch, Multiple solutions of steady MHD flow of dilatant fluids, *Eur. J. Pure Appl. Math.* 1 (2) (2008) 11–20.
- [14] X. Hu, Z. Dong, F. Huang, Y. Chen, Symmetry reductions and exact solutions of the (2+1)-dimensional Navier–Stokes equations, *Z. Nat.forsch. A* 65 (2010) 504–510.
- [15] M.Y. Hussaini, W.D. Lakin, Existence and non-uniqueness of similarity solutions of a boundary-layer problem, *Q. J. Mech. Appl. Math.* 39 (1986) 177–191.
- [16] O.A. Ladyzhenskaya, *Boundary Value Problems of Mathematical Physics*, American Mathematical Society, Providence, RI, 1970.
- [17] J.H. Merkin, On dual solutions occurring in mixed convection in a porous medium, *J. Eng. Math.* 20 (1985) 171–179.
- [18] H. Schlichting, *Boundary Layer Theory*, McGraw-Hill Press, New York, 1979.
- [19] L. Sedov, *Similarity and Dimensional Methods in Mechanics*, CRC Press, Boca Raton, 1993.
- [20] D. Wei, S. Al-Ashhab, Similarity solutions for a non-newtonian power-law fluid flow, *Appl. Math. Mech. (English Ed.)* 35 (2014) 1155–1166, <http://dx.doi.org/10.1007/s10483-014-1854-6>.
- [21] L. Zheng, X. Zhang, J. He, Existence and estimate of positive solutions to a nonlinear singular boundary value problem in the theory of dilatant non-Newtonian fluids, *Math. Comput. Modelling* 45 (2007) 387–393.